

Gene therapy with HSV encoding p55TNFR gene for HIV neuropathic pain: an evidence-based mini-review

Hirotsugu Kanda^{1,2}, Shue Liu¹, Megumi Kanao^{1,2}, Hyun Yi¹, Takafumi Iida^{1,2}, Wan Huang¹, Takayuki Kunisawa², David A. Lubarsky¹ and Shuanglin Hao^{1*}

¹Department of Anesthesiology, University of Miami Miller School of Medicine, Miami, Florida 33136

²Department of Anesthesiology and Critical Care Medicine, Asahikawa Medical University, Japan 078-8510

*The work was partially reported at the 2016 annual meeting of American Society of Anesthesiologists in Chicago.

Abstract

While effective antiretroviral treatment makes human immunodeficiency virus (HIV)-related death decreased dramatically, neuropathic pain becomes one of the most common complications in patients with HIV/acquired immunodeficiency syndrome (AIDS). The exact mechanisms of HIV-related neuropathic pain are not well understood yet, and no effective therapy is for HIV-pain. Evidence has shown that proinflammatory factors (e.g., tumor necrosis factor alpha (TNF α)) released from glia, are critical to contributing to chronic pain. Preclinical studies have demonstrated that non-replicating herpes simplex virus (HSV)-based vector expressing human enkephalin reduces inflammatory pain, neuropathic pain, or cancer pain in animal models. In this review, we describe recent advances in the use of HSV-based gene transfer for the treatment of HIV pain, with a special focus on the use of HSV-mediated soluble TNF receptor I (neutralizing TNF α in function) in HIV neuropathic pain model.

Keywords

HIV, Neuropathic pain, Soluble TNF receptor, and Gene therapy

our work and others focusing on the pathophysiological mechanisms of HIV-neuropathic pain and gene therapy.

HIV-Related Neuropathic Pain

HIV sensory neuropathies contain distal sensory polyneuropathy as results of both HIV infection and antiretroviral drug-induced toxic neuropathies [3-6]. Clinical characteristics of distal sensory polyneuropathy and ART-induced toxic neuropathies are very similar. Neurotoxic ART has even been removed from pharmacies entirely in developed countries. Evidence shows that many people with HIV alive today, have ever been on numerous therapeutic regimens with neurotoxic drugs, and that they have already developed persistent painful neuropathy [7,8]. HIV-NP is typically bilateral, of gradual onset, and described as 'aching', 'painful numbness', or 'burning' [9]. Pathological feature of HIV-NP includes loss of sensory neurons of the dorsal root ganglion (DRG), Wallerian degeneration of the long axons in distal area, infiltration of macrophage into the DRG, and a 'dying back' sensory neuropathy [10-14]. However, the precise mechanisms of HIV-NP remain unknown yet and no effective therapy for HIV-NP.

Proinflammatory factors

Early studies have demonstrated that glia infected/activated by HIV release proinflammatory factors, such as tumor necrosis factor alpha (TNF α) and interleukin 1 (IL-1) [15]. Infiltration of inflammatory lymphocyte and macrophage to the DRG of AIDS patients produces pro-inflammatory cytokines including TNF α [12,16-20]. There is an increased TNF α in human CSF [21-25] and brain tissue [25-28] in patients with HIV. An interaction of TNF α and HIV infection enhances toxic chemokine products [29,30]. It is known that proinflammatory cytokines play an important role in the development and maintenance of neuropathic pain [31-35]. Proinflammatory mediators are critical to enhancing HIV-NP [36]. Intrathecal administration of gp120 induces acute pain and spinal proinflammatory cytokine release [37]. Pe-

Introduction

The United States Centers for Disease Control reports that an estimated 1.1 million people were living with human immunodeficiency virus (HIV) at the end of 2014 and that 39,513 people in 2015 were diagnosed with HIV infection in the United States (<https://www.cdc.gov/hiv/basics/statistics.html>, last date accessed June 24, 2017). Although effective antiretroviral therapy (ART) makes HIV become a treatable, chronic disease [1,2], new challenges are emerging in managing HIV. Chronic pain becomes one of the most common complications in patients with HIV/acquired immunodeficiency syndrome (AIDS). HIV-related pain is often underestimated in HIV/AIDS patients while the main focus is on immunosuppression and opportunistic infections. HIV neuropathic pain (HIV-NP) is refractory, and the current available chronic pain therapies are not effective to HIV-NP. This article reviews current researches from

peripheral gp120 increases TNF α within the nerve trunk [38], intense glial activation in the spinal cord in parallel with neuropathic pain behaviors [38]. We have reported that peripheral gp120 application onto the rat sciatic nerve upregulates TNF α in the L4/5 DRGs and spinal cord [39]. Systemic 2',3'-dideoxycytidine (ddC), one drug of ART lowers mechanical threshold [40,41] and increases both mRNA and protein of TNF α in the spinal cord dorsal horn (SCDH) [41]. Inhibition of TNF α or soluble TNF receptor reduces mechanical allodynia induced by gp120 application [41].

Therefore, it is possible that TNF α signal is involved in the induction and/or progression of HIV-NP.

Reactive oxygen species and C/EBP β in HIV

Oxidative stress evokes many signaling events [42]. Mitochondria are the main source of reactive oxygen species (ROS). ROS plays a role in different pain models [43-48]. ROS scavengers produce a strong antinociceptive effect in persistent pain models [49]. Oxidative stress is involved in the pathogenesis of neuroAIDS [50]. HIV infection and ART can evoke rapid neurotoxicity [51]. Either HIV gp120 or ddC plays a role in initiation and/or intensification of ROS [52,53]. Intrathecal gp120 induces spinal release of nitric oxide (NO) as well as proinflammatory cytokines; pretreatment with NO synthase (NOS) inhibitor abolishes gp120-induced mechanical allodynia [54]. Importantly, ROS evoked by HIV infection, induces apoptosis through TNF α and its receptors [52]. Mitochondrial DNA (mtDNA) is critical for oxidative phosphorylation complex I proteins. DNA polymerase- γ is important for replication of mtDNA. ARTs inhibit Polymerase- γ , resulting in mitochondrial respiratory chain dysfunction and oxidative phosphorylation deficits [51]. Systemic ddC induces neuropathic pain and lowers the activity of endogenous manganese superoxide dismutase (SOD2) in the SCDH; ROS scavengers significantly reduce mechanical allodynia [55].

CCAAT/enhancer binding proteins (C/EBPs) are transcriptional factors in cell development and induction of inflammatory factors in the peripheral and central nervous system [56]. C/EBP β plays an important role in a variety of HIV disease stages [57]. An increase in C/EBP β mRNA is found in the brain tissue of HIV-1 encephalitis patients [58]. We have found that combination of peripheral gp120 with systemic ddC increases pC/EBP β in the SCDH [59], suggesting that pC/EBP β plays a role in HIV-NP.

HSV Vector for Gene Therapy of Neuropathic Pain

During natural infection of herpes simplex virus (HSV), HSV is carried by retrograde axonal transport from the site of original inoculation to the neuronal perikaryon. Latently infected neurons function normally and are

not rejected by the host immune response (see review [60,61]). HSV-1 genome is a linear double-stranded DNA, and has more than 75 genes coded in the 152 kb genome [60,61]. HSV genes are expressed in a well-ordered temporal cascade of immediate early (IE) genes, followed by early genes, and subsequently late gene products; both early genes and the late genes require synthesis of IE gene products [60,61]. Deleting essential IE genes from the HSV genome makes it non-replicating recombinant [62], but the virus are still able to be used to effectively deliver target gene products [63,64]. Gene transfer mediated by HSV vector may provide a promising approach to the management of neuropathic pain. HSV vector encoding human preproenkephalin gene after transduction of DRG neurons by hindpaw injection [65], produces an antinociceptive effect in different pain models [66-68]. We have reported that HSV vectors expressing enkephalin, p55 TNF soluble receptor (p55TNFSR), interleukin-10, and interleukin-4 produce antinociceptive effects in preclinical pain models [69-75]. Fink and colleagues reported phase 1 clinical trial using HSV vector encoding human preproenkephalin in patients with cancer pain [76]. The clinical trial assessed the safety and explored the potential efficacy of this approach in humans, indicating that it may be effective in reducing cancer pain [77].

The distribution of systemically administered drugs to the brain may be limited by the blood-brain barrier [65], and they produce systemic side effects. Gene transfer that permanently release gene products, might be a useful alternative to regular pharmacological approaches [65]. Gene transfer of HSV vector may represent a platform technology---nerve targeting drug delivery system [77]. Viral vectors, however, show toxicity and inflammation from 'leaky' expression of viral genes and reaction to the vector coat protein in pre-immune animals [65,78]. Despite these limitations, ours and other studies have shown that HSV vector is still a highly effective gene delivery approach to treating peripheral and central nervous diseases [65,79,80].

TNFSR mediated by HSV vector produces antiallodynic effect in HIV-NP

Our report has shown that the HIV gp120 application onto the sciatic nerve induces upregulation of TNF α , C-X-C chemokine receptor type 4 (CXCR4, a co-receptor of HIV), stromal cell-derived factor 1- α (SDF1- α , CXCR4 ligand) in both the DRG and the lumbar spinal dorsal horn [81]. Soluble TNF receptor (TNFSR) blocks bioactivity of TNF α . HSV vector encoding p55TNFSR gene (TOTNFSR) reduces mechanical allodynia and lowers TNF α , CXCR4 and SDF1- α induced by gp120 in the DRG and SCDH [81], suggesting that the pathway of TNF α to the CXCR4/SDF1 has an important role in the HIV-NP and that inhibiting proinflammatory cytokines/chemokines reduce

neuropathic pain. In another model of HIV-NP induced by intraperitoneal ddC [40], ddC induces upregulation of TNF α , SDF1- α , and CXCR4 in both the lumbar spinal

cord and the L4/5 DRG; T0TNFSR reduced mechanical allodynia and suppressed TNF α , SDF1- α , and CXCR4 in the lumbar SCDH and DRG [82], indicating that TNF α is




Figure 1: Antinociceptive effect of p55TNFRI mediated by HSV vectors. Mechanical sensitivity was examined through the measurement of foot withdrawal frequencies to a sequential series of calibrated von Frey filaments applied in ascending order to the plantar surface of the foot (98). HSV vector T0TNFSR or T0Z was inoculated into the hindpaws 1 week post gp120/ddC. The occurrence of foot withdrawal for each trial was expressed as a percentage response frequency. Two weeks after HSV vectors, foot withdrawal frequencies to calibrated von Frey filaments in rats with subcutaneous inoculation of T0TNFSR were significantly lower than that in T0Z at filaments of 3.6, 5.5, 8.5, and 11.8 gram, * $P < 0.05$, *** $P < 0.001$ vs. T0TNFSR, t test, $n = 6$.




Figure 2: The expression of p55TNFRI mediated by the HSV vectors. One week after gp120/ddC, T0TNFSR or T0Z was inoculated into the hindpaws. On day 14 post HSV vector, the L4/5 dorsal root ganglion (DRG) was harvested, and western blot assays were conducted for testing TNFRI. T0TNFSR injection significantly induced the expression of TNFRI compared with T0Z in the L4/5 DRG, ** $P < 0.01$ vs. T0Z, t test, $n = 6$.




Figure 3: The effect of p55TNFRI mediated by the HSV vectors on mitochondrial superoxide in the SCDH at 2 weeks post HSV vectors. One week post gp120/ddC, neuropathic rats received hindpaw injection of T0TNFSR or T0Z into ipsilateral hindpaw of rats. Two weeks after HSV vector, MitoSox Red was intrathecally injected 70 min prior to perfusion. The representative image of MitoSox red for mitochondrial superoxide in sham+T0Z, gp120/ddC+T0Z, and gp120/ddC+T0TNFSR, was shown in Figure A, B, and C, respectively, scale bar, 50 μ m. (D) The number of mitochondrial superoxide positive cells in the SCDH lamina I-II and III-V was shown, ** $P < 0.01$, *** $P < 0.001$, one way ANOVA with *post hoc* PLSD test, mean \pm SEM, $n = 5-6$ rats.

involved in the ARTs-related pain through the SDF1- α /CXCR4 system.

We have reported that combination of peripheral gp120 with systemic ddC (gp120/ddC) lowers mechanical threshold for more than 3 weeks, and that the minimum of mechanical threshold occurs around 2 weeks after gp120/ddC [59,83,84]. Previous studies show that HSV vector T0TNFSR reduces neuropathic pain induced by spinal nerve injury [69]. In gp120/ddC model, 2 weeks post HSV vector, T0TNFSR significantly reduced foot withdrawal frequencies (Figure 1), and increased the expression of soluble TNFRI in the L4/5 DRG (Figure 2).

TNFSR mediated by the HSV vector reduces mitochondrial superoxide in gp120/ddC model

Oxidative stress causes many signaling events [42]. HIV gp120 or ddC induces ROS [52,53]. HIV gp120 application onto the sciatic nerve upregulates spinal mitochondrial superoxide [73,85]. We reported that gp120/ddC increased spinal mitochondrial superoxide [59,84] using MitoSox positive cell imaging (a marker of mitochondrial superoxide) [86,87]. Figure 3A-C showed the representative Mi-

toSox positive cell imaging in the gp120/ddC model. The increased number of MitoSox positive cells in the gp120/ddC model was decreased by HSV vector T0TNFSR (Figure 3D), suggesting that TNFSR suppresses neuropathic pain through reducing spinal ROS.

TNFSR mediated by the HSV vector inhibits pC/EBP β in the gp120/ddC neuropathic pain model

C/EBP plays a role in induction of inflammatory mediators in CNS [56]. HIV patients show upregulation of C/EBP β mRNA in the brain tissue [58]. We have shown that HIV-NP increases phosphorylation of C/EBP β (pC/EBP β) [59]. Figure 4A-C revealed the representative pC/EBP β -IR images in the gp120/ddC model. Treatment with gp120/ddC increased pC/EBP β -IR expression; the upregulated pC/EBP β -IR was suppressed by T0TNFSR (Figure 4D), suggesting that TNFSR reduces neuropathic pain through decreasing spinal pC/EBP β .

The relationship of TNF α /TNF receptor activity and ROS or C/EBP β in HIV-NP is still not clear. HIV gp120/ddC induces release of TNF α [83]. Through TNF receptor, TNF α triggers a cascade of events [88]. TNF α activates NMDA




Figure 4: The effect of p55TNFSR mediated by the HSV vectors on pC/EBP β in the SDH at 2 weeks post HSV vectors. One week post gp120/ddC, neuropathic rats received injection of T0TNFSR or T0Z into ipsilateral hindpaw of rats. Two weeks after HSV vector, spinal pC/EBP β -immunoreactivity (pC/EBP β -IR) was examined using immunohistochemistry. The representative images of pC/EBP β -IR in sham+T0Z, gp120/ddC+T0Z, and gp120/ddC+T0TNFSR were shown in Figure A, B, and C, respectively, scale bar, 50 μ m. (D) The quantitative signals of pC/EBP β -IR in the SCDH were shown, ** $P < 0.01$, one way ANOVA with post hoc PLSD test, mean \pm SEM, $n=6$ rats.

receptors to increase Ca^{2+} influx [89]. Indeed, HIV gp120 increases intracellular free Ca^{2+} concentration in the mice SCDH cells [90]. ddC also increases spinal cytosolic Ca^{2+} concentration in painful neuropathy [91]. There is an interplay between cytosolic Ca^{2+} and mitochondrial ROS [92-94]. Spinal pCREB may make a contribution to the development of chronic pain [95]. Cytosolic Ca^{2+} may induce transcriptional factor CREB regulating C/EBP β activity [96]. CREB binds C/EBP β gene promoter, inducing the endogenous C/EBP β expression [97]. Therefore, it is possible that TNF α -TNFR induces ROS, or pCREB/ C/EBP β in HIV-NP, which need to be examined in the near future.

In summary, glia infected or activated after HIV release proinflammatory factors, such as TNF α . TNF α -TNF receptor signal may induce ROS or C/EBP β in HIV-NP through complex pathway in the model of HIV-NP. Gene transfer using the HSV vector encoding the gene of TNF soluble receptor reduced neuropathic pain in animal studies, providing additional potential approach for successful treatment of HIV neuropathic pain.

Conflict of Interest

All authors declare no conflict of interest in the work.

Acknowledgements

Supported by grants from the National Institutes of Health DA026734 (SH), DA025527 (SH), NS066792 (SH) and DA34749 (SH), and by the Department of Anesthesiology, the University of Miami Miller School of Medicine, Miami, FL.

Corresponding Author: Shuanglin Hao, MD, PhD, Professor & Director for Preclinical and Basic Research, Department of Anesthesiology, University of Miami Miller School of Medicine Miami, FL33136, USA; Email: shao@med.miami.edu

Editor: Yuan-Xiang Tao, Ph.D., M.D., Professor and Vice Chair, Director, Center for Pain Medicine Research Department of Anesthesiology, Editor in Chief, The Translational Perioperative and Pain Medicine, Rutgers, The State University of New Jersey, New Jersey Medical School, 185 S. Orange Ave., MSB, E-661 Newark, NJ 07103. Phone: 973-972-9812 (O) Email: yt211@njms.rutgers.edu

Additional publication details

Journal short name: Transl Perioper & Pain Med

Received Date: August 12, 2017

Accepted Date: September 21, 2017

Published Date: October 03, 2017

Citation: Kanda H, Liu S, Kanao M, Yi H, Iida T, et al. Gene therapy with HSV encoding p55TNFR gene for HIV neuropathic pain: an evidence-based minireview. Transl Perioper & Pain Med 2017; 4 (4):24-32

Copyright: © 2017 Kanda H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

References

- Colvin CJ. HIV/AIDS, chronic diseases and globalisation. *Globalization and health* 2011;7:31.
- Deeks SG, Lewin SR, Havlir DV. The end of AIDS: HIV infection as a chronic disease. *Lancet* 2013;382:1525-33.
- Keswani SC, Pardo CA, Cherry CL, Hoke A, McArthur JC. HIV-associated sensory neuropathies. *Aids* 2002;16:2105-17.
- Nath A. Neurologic Complications of Human Immunodeficiency Virus Infection. *Continuum (Minneapolis)* 2015;21:1557-76.
- Hoke A, Cornblath DR. Peripheral neuropathies in human immunodeficiency virus infection. *Suppl Clin Neurophysiol* 2004;57:195-210.
- Cherry CL, Wadley AL, Kamerman PR. Painful HIV-associated sensory neuropathy. *Pain Manag* 2012;2:543-52.
- Kranick SM, Nath A. Neurologic complications of HIV-1 infection and its treatment in the era of antiretroviral therapy. *Continuum (Minneapolis)* 2012;18:1319-37.
- Leger PD, Johnson DH, Robbins GK, Shafer RW, Clifford DB, Li J, McLaren PJ, Haas DW. Genome-wide association study of peripheral neuropathy with D-drug-containing regimens in AIDS Clinical Trials Group protocol 384. *J Neurovirol* 2014;20:304-8.
- Cornblath DR, McArthur JC. Predominantly sensory neuropathy in patients with AIDS and AIDS-related complex. *Neurology* 1988;38:794-6.
- Wulff EA, Wang AK, Simpson DM. HIV-associated peripheral neuropathy: epidemiology, pathophysiology and treatment. *Drugs* 2000;59:1251-60.
- Keswani SC, Polley M, Pardo CA, Griffin JW, McArthur JC, Hoke A. Schwann cell chemokine receptors mediate HIV-1 gp120 toxicity to sensory neurons. *Ann Neurol* 2003;54:287-96.
- Pardo CA, McArthur JC, Griffin JW. HIV neuropathy: insights in the pathology of HIV peripheral nerve disease. *J Peripher Nerv Syst* 2001;6:21-7.
- McCarthy BG, Hsieh ST, Stocks A, Hauer P, Macko C, Cornblath DR, Griffin JW, McArthur JC. Cutaneous innervation in sensory neuropathies: evaluation by skin biopsy. *Neurology* 1995;45:1848-55.
- Verma A. Epidemiology and clinical features of HIV-1 associated neuropathies. *J Peripher Nerv Syst* 2001;6:8-13.
- Merrill JE, Chen IS. HIV-1, macrophages, glial cells, and cytokines in AIDS nervous system disease. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 1991;5:2391-7.
- Nagano I, Shapshak P, Yoshioka M, Xin K, Nakamura S, Bradley WG. Increased NADPH-diaphorase reactivity and cytokine expression in dorsal root ganglia in acquired immunodeficiency syndrome. *Journal of the neurological sciences* 1996;136:117-28.
- Rizzuto N, Cavallaro T, Monaco S, Morbin M, Bonetti B, Ferrari S, Galiazzo-Rizzuto S, Zanette G, Bertolasi L. Role of HIV in the pathogenesis of distal symmetrical peripheral neuropathy. *Acta Neuropathol* 1995;90:244-50.
- Shapshak P, Nagano I, Xin K, Bradley W, McCoy CB, Sun NC, Stewart RV, Yoshioka M, Petito C, Goodkin K, et al. HIV-1 heterogeneity and cytokines. *Neuropathogenesis. Advances in experimental medicine and biology* 1995;373:225-38.
- Yoshioka M, Shapshak P, Srivastava AK, Stewart RV, Nelson SJ, Bradley WG, Berger JR, Rhodes RH, Sun NC, Nakamura S. Expression of HIV-1 and interleukin-6 in lumbosacral dorsal root ganglia of patients with AIDS. *Neurology* 1994;44:1120-30.
- Wesselingh SL, Glass J, McArthur JC, Griffin JW, Griffin DE. Cytokine dysregulation in HIV-associated neurological disease. *Adv Neuroimmunol* 1994;4:199-206.
- Grimaldi LM, Martino GV, Franciotta DM, Brustia R, Castagna A, Pristera R, Lazzarin A. Elevated alpha-tumor necrosis factor levels in spinal fluid from HIV-1-infected patients with central nervous system involvement. *Ann Neurol* 1991;29:21-5.
- Mastroianni CM, Paoletti F, Valenti C, Vullo V, Jirillo E, Delia S. Tumour necrosis factor (TNF-alpha) and neurological disorders in HIV infection. *J Neurol Neurosurg Psychiatry* 1992;55:219-21.
- Perrella O, Carrieri PB, Guarnaccia D, Soscia M. Cerebrospinal fluid cytokines in AIDS dementia complex. *J Neurol* 1992;239:387-8.
- Ciardi M, Sharief MK, Thompson EJ, Salotti A, Vullo V, Sorice F, Cirelli A. High cerebrospinal fluid and serum levels of tumor necrosis factor-alpha in asymptomatic HIV-1 seropositive individuals. Correlation with interleukin-2 and soluble IL-2 receptor. *Journal of the neurological sciences* 1994;125:175-9.
- Tyor WR, Glass JD, Griffin JW, Becker PS, McArthur JC, Bezman L, Griffin DE. Cytokine expression in the brain during the acquired immunodeficiency syndrome. *Ann Neurol* 1992;31:349-60.
- Achim CL, Heyes MP, Wiley CA. Quantitation of human immunodeficiency virus, immune activation factors, and quinolinic acid in AIDS brains. *J Clin Invest* 1993;91:2769-75.
- An SF, Ciardi A, Giometto B, Scaravilli T, Gray F, Scaravilli F. Investigation on the expression of major histocompatibility complex class II and cytokines and detection of HIV-1 DNA within brains of asymptomatic and symptomatic HIV-1-positive patients. *Acta Neuropathol* 1996;91:494-503.
- Vitkovic L, da Cunha A, Tyor WR. Cytokine expression and pathogenesis in AIDS brain. *Res Publ Assoc Res Nerv Ment Dis* 1994;72:203-22.
- Williams R, Dhillon NK, Hegde ST, Yao H, Peng F, Callen S, Chebloune Y, Davis RL, Buch SJ. Proinflammatory cytokines and HIV-1 synergistically enhance CXCL10 expression in human astrocytes. *Glia* 2009;57:734-43.
- Kaul M, Garden GA, Lipton SA. Pathways to neuronal injury and apoptosis in HIV-associated dementia. *Nature* 2001;410:988-94.
- Raghavendra V, Rutkowski MD, DeLeo JA. The role of spinal neuroimmune activation in morphine tolerance/hyperalgesia in neuropathic and sham-operated rats. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2002;22:9980-9.
- Watkins LR, Milligan ED, Maier SF. Glial activation: a driving force for pathological pain. *Trends in neurosciences* 2001;24:450-5.

33. DeLeo JA, Tanga FY, Tawfik VL. Neuroimmune activation and neuroinflammation in chronic pain and opioid tolerance/hyperalgesia. *Neuroscientist* 2004;10:40-52.
34. Leung L, Cahill CM. TNF-alpha and neuropathic pain--a review. *J Neuroinflammation* 2010;7:27.
35. Gao YJ, Zhang L, Ji RR. Spinal injection of TNF-alpha-activated astrocytes produces persistent pain symptom mechanical allodynia by releasing monocyte chemoattractant protein-1. *Glia* 2010;58:1871-80.
36. Hao S. The Molecular and Pharmacological Mechanisms of HIV-Related Neuropathic Pain. *Current Neuropharmacology* 2013;11:499-512.
37. Milligan ED, O'Connor KA, Nguyen KT, Armstrong CB, Twining C, Gaykema RP, Holguin A, Martin D, Maier SF, Watkins LR. Intrathecal HIV-1 envelope glycoprotein gp120 induces enhanced pain states mediated by spinal cord proinflammatory cytokines. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2001;21:2808-19.
38. Herzberg U, Sagen J. Peripheral nerve exposure to HIV viral envelope protein gp120 induces neuropathic pain and spinal gliosis. *J Neuroimmunol* 2001;116:29-39.
39. Zheng W, Ouyang H, Zheng X, Liu S, Mata M, Fink DJ, Hao S. Glial TNFalpha in the spinal cord regulates neuropathic pain induced by HIV gp120 application in rats. *Mol Pain* 2011;7:40.
40. Joseph EK, Chen X, Khasar SG, Levine JD. Novel mechanism of enhanced nociception in a model of AIDS therapy-induced painful peripheral neuropathy in the rat. *Pain* 2004;107:147-58.
41. Zheng X, Ouyang H, Liu S, Mata M, Fink DJ, Hao S. TNFalpha is involved in neuropathic pain induced by nucleoside reverse transcriptase inhibitor in rats. *Brain Behav Immun* 2011;25:1668-76.
42. Chandra J, Samali A, Orrenius S. Triggering and modulation of apoptosis by oxidative stress. *Free Radic Biol Med* 2000;29:323-33.
43. Crisp T, Minus TO, Coleman ML, Giles JR, Cibula C, Finnerly EP. Aging, peripheral nerve injury and nociception: effects of the antioxidant 16-desmethyltirilazad. *Behav Brain Res* 2006;166:159-65.
44. Khalil Z, Liu T, Helme RD. Free radicals contribute to the reduction in peripheral vascular responses and the maintenance of thermal hyperalgesia in rats with chronic constriction injury. *Pain* 1999;79:31-7.
45. Liu D, Liu J, Sun D, Wen J. The time course of hydroxyl radical formation following spinal cord injury: the possible role of the iron-catalyzed Haber-Weiss reaction. *J Neurotrauma* 2004;21:805-16.
46. Kim HK, Park SK, Zhou JL, Taglialatela G, Chung K, Coggeshall RE, Chung JM. Reactive oxygen species (ROS) play an important role in a rat model of neuropathic pain. *Pain* 2004;111:116-24.
47. Mao YF, Yan N, Xu H, Sun JH, Xiong YC, Deng XM. Edaravone, a free radical scavenger, is effective on neuropathic pain in rats. *Brain Res* 2009;1248:68-75.
48. Schwartz ES, Kim HY, Wang J, Lee I, Klann E, Chung JM, Chung K. Persistent pain is dependent on spinal mitochondrial antioxidant levels. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2009;29:159-68.
49. Kim HY, Chung JM, Chung K. Increased production of mitochondrial superoxide in the spinal cord induces pain behaviors in mice: The effect of mitochondrial electron transport complex inhibitors. *Neurosci Lett* 2008;447:87-91.
50. Mollace V, Nottet HS, Clayette P, Turco MC, Muscoli C, Salvemini D, Perno CF. Oxidative stress and neuroAIDS: triggers, modulators and novel antioxidants. *Trends in neurosciences* 2001;24:411-6.
51. Yamaguchi T, Katoh I, Kurata S. Azidothymidine causes functional and structural destruction of mitochondria, glutathione deficiency and HIV-1 promoter sensitization. *Eur J Biochem* 2002;269:2782-8.
52. Perl A, Banki K. Genetic and metabolic control of the mitochondrial transmembrane potential and reactive oxygen intermediate production in HIV disease. *Antioxid Redox Signal* 2000;2:551-73.
53. Lewis W, Day BJ, Copeland WC. Mitochondrial toxicity of NRTI antiviral drugs: an integrated cellular perspective. *Nat Rev Drug Discov* 2003;2:812-22.
54. Holguin A, O'Connor KA, Biedenkapp J, Campisi J, Wieseler-Frank J, Milligan ED, Hansen MK, Spataro L, Maksimova E, Bravmann C, Martin D, Fleshner M, Maier SF, Watkins LR. HIV-1 gp120 stimulates proinflammatory cytokine-mediated pain facilitation via activation of nitric oxide synthase-I (nNOS). *Pain* 2004;110:517-30.
55. Zheng W, Zheng X, Liu S, Hao S. ROS in the spinal cord is involved in the neuropathic pain induced by antiretroviral drug in rats Society for Neuroscience annual meeting. Washington, DC, 2011.
56. Alberini CM. Transcription factors in long-term memory and synaptic plasticity. *Physiol Rev* 2009;89:121-45.
57. Huber R, Pietsch D, Panterodt T, Brand K. Regulation of C/EBPbeta and resulting functions in cells of the monocytic lineage. *Cell Signal* 2012;24:1287-96.
58. Fields J, Gardner-Mercer J, Borgmann K, Clark I, Ghorpade A. CCAAT/enhancer binding protein beta expression is increased in the brain during HIV-1-infection and contributes to regulation of astrocyte tissue inhibitor of metalloproteinase-1. *J Neurochem* 2011;118:93-104.
59. Kanao M, Kanda H, Huang W, Liu S, Yi H, Candiotti KA, Lubarsky DA, Levitt RC, Hao S. Gene Transfer of Glutamic Acid Decarboxylase 67 by Herpes Simplex Virus Vectors Suppresses Neuropathic Pain Induced by Human Immunodeficiency Virus gp120 Combined with ddC in Rats. *Anesth Analg* 2015;120:1394-404.
60. Fink DJ, DeLuca NA, Goins WF, Glorioso JC. Gene transfer to neurons using herpes simplex virus-based vectors. *Annu Rev Neurosci* 1996;19:265-87.
61. Manservigi R, Argnani R, Marconi P. HSV Recombinant Vectors for Gene Therapy. *Open Virol J* 2010;4:123-56.
62. DeLuca NA, McCarthy AM, Schaffer PA. Isolation and characterization of deletion mutants of herpes simplex virus type 1 in the gene encoding immediate-early regulatory protein ICP4. *J Virol* 1985;56:558-70.
63. Mata M, Hao S, Fink DJ. Applications of gene therapy to the treatment of chronic pain. *Curr Gene Ther* 2008;8:42-8.
64. Glorioso JC, Fink DJ. Herpes vector-mediated gene transfer in the treatment of chronic pain. *Mol Ther* 2009;17:13-8.

65. Simonato M, Bennett J, Boulis NM, Castro MG, Fink DJ, Goins WF, Gray SJ, Lowenstein PR, Vandenberghe LH, Wilson TJ, Wolfe JH, Glorioso JC. Progress in gene therapy for neurological disorders. *Nat Rev Neurol* 2013;9:277-91.
66. Wilson SP, Yeomans DC, Bender MA, Lu Y, Goins WF, Glorioso JC. Antihyperalgesic effects of infection with a preproenkephalin-encoding herpes virus. *Proceedings of the National Academy of Sciences of the United States of America* 1999;96:3211-6.
67. Goss JR, Mata M, Goins WF, Wu HH, Glorioso JC, Fink DJ. Antinociceptive effect of a genomic herpes simplex virus-based vector expressing human proenkephalin in rat dorsal root ganglion. *Gene Ther* 2001;8:551-6.
68. Braz J, Beaufour C, Coutaux A, Epstein AL, Cesselin F, Hamon M, Pohl M. Therapeutic efficacy in experimental polyarthritis of viral-driven enkephalin overproduction in sensory neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2001;21:7881-8.
69. Hao S, Mata M, Glorioso JC, Fink DJ. Gene transfer to interfere with TNFalpha signaling in neuropathic pain. *Gene Ther* 2007;14:1010-6.
70. Hao S, Mata M, Glorioso JC, Fink DJ. HSV-mediated expression of interleukin-4 in dorsal root ganglion neurons reduces neuropathic pain. *Mol Pain* 2006;2:6.
71. Zhou Z, Peng X, Hao S, Fink DJ, Mata M. HSV-mediated transfer of interleukin-10 reduces inflammatory pain through modulation of membrane tumor necrosis factor alpha in spinal cord microglia. *Gene Ther* 2008;15:183-90.
72. Hao S, Mata M, Goins W, Glorioso JC, Fink DJ. Transgene-mediated enkephalin release enhances the effect of morphine and evades tolerance to produce a sustained antiallodynic effect in neuropathic pain. *Pain* 2003;102:135-42.
73. Kanda H, Kanao M, Liu S, Yi H, Iida T, Levitt RC, Candiotti KA, Lubarsky DA, Hao S. HSV vector-mediated GAD67 suppresses neuropathic pain induced by perineural HIV gp120 in rats through inhibition of ROS and Wnt5a. *Gene Ther* 2016;23:340-8.
74. Majima T, Funahashi Y, Takai S, Goins WF, Gotoh M, Tyagi P, Glorioso JC, Yoshimura N. Herpes Simplex Virus Vector-Mediated Gene Delivery of Poreless TRPV1 Channels Reduces Bladder Overactivity and Nociception in Rats. *Human gene therapy* 2015;26:734-42.
75. Oguchi T, Funahashi Y, Yokoyama H, Nishizawa O, Goins WF, Goss JR, Glorioso JC, Yoshimura N. Effect of herpes simplex virus vector-mediated interleukin-4 gene therapy on bladder overactivity and nociception. *Gene Ther* 2013;20:194-200.
76. Wolfe D, Mata M, Fink DJ. A human trial of HSV-mediated gene transfer for the treatment of chronic pain. *Gene Ther* 2009;16:455-60.
77. Fink DJ, Wechuck J, Mata M, Glorioso JC, Goss J, Krisky D, Wolfe D. Gene therapy for pain: results of a phase I clinical trial. *Ann Neurol* 2011;70:207-12.
78. Nayak S, Herzog RW. Progress and prospects: immune responses to viral vectors. *Gene Ther* 2010;17:295-304.
79. Guedon JM, Wu S, Zheng X, Churchill CC, Glorioso JC, Liu CH, Liu S, Vulchanova L, Bekker A, Tao YX, Kinchington PR, Goins WF, Fairbanks CA, Hao S. Current gene therapy using viral vectors for chronic pain. *Mol Pain* 2015;11:27.
80. Hao S, Liu S, Zheng X, Zheng W, Ouyang H, Mata M, Fink DJ. The role of TNFalpha in the periaqueductal gray during naloxone-precipitated morphine withdrawal in rats. *Neuropsychopharmacology* 2011;36:664-76.
81. Huang W, Zheng W, Liu S, Zeng W, Levitt RC, Candiotti KA, Lubarsky DA, Hao S. HSV-mediated p55TNFSR reduces neuropathic pain induced by HIV gp120 in rats through CXCR4 activity. *Gene Ther* 2014;21:328-36.
82. Huang W, Zheng W, Ouyang H, Yi H, Liu S, Zeng W, Levitt RC, Candiotti KA, Lubarsky DA, Hao S. Mechanical allodynia induced by nucleoside reverse transcriptase inhibitor is suppressed by p55TNFSR mediated by herpes simplex virus vector through the SDF1alpha/CXCR4 system in rats. *Anesth Analg* 2014;118:671-80.
83. Zheng W, Huang W, Liu S, Levitt RC, Candiotti KA, Lubarsky DA, Hao S. IL-10 mediated by herpes simplex virus vector reduces neuropathic pain induced by HIV gp120 combined with ddC in rats. *Mol Pain* 2014;10:49.
84. Iida T, Yi H, Liu S, Huang W, Kanda H, Lubarsky DA, Hao S. Spinal CPEB-mtROS-CBP signaling pathway contributes to perineural HIV gp120 with ddC-related neuropathic pain in rats. *Experimental neurology* 2016;281:17-27.
85. Kanda H, Liu S, Iida T, Yi H, Huang W, Levitt R, Lubarsky DA, Candiotti KA, Hao S. Inhibition of mitochondrial fission protein reduced mechanical allodynia and suppressed spinal mitochondrial superoxide induced by perineural HIV gp120 in rats. *Anesth Analg* 2016;122:264-72.
86. Kim HY, Lee KY, Lu Y, Wang J, Cui L, Kim SJ, Chung JM, Chung K. Mitochondrial Ca²⁺ uptake is essential for synaptic plasticity in pain. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2011;31:12982-91.
87. Schwartz ES, Lee I, Chung K, Chung JM. Oxidative stress in the spinal cord is an important contributor in capsaicin-induced mechanical secondary hyperalgesia in mice. *Pain* 2008;138:514-24.
88. Ferrari LF, Bogen O, Levine JD. Role of nociceptor alpha-CaMKII in transition from acute to chronic pain (hyperalgesic priming) in male and female rats. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 2013;33:11002-11.
89. Ohno M, Frankland PW, Silva AJ. A pharmacogenetic inducible approach to the study of NMDA/alphaCaMKII signaling in synaptic plasticity. *Curr Biol* 2002;12:654-6.
90. Minami T, Matsumura S, Mabuchi T, Kobayashi T, Sugimoto Y, Ushikubi F, Ichikawa A, Narumiya S, Ito S. Functional evidence for interaction between prostaglandin EP3 and kappa-opioid receptor pathways in tactile pain induced by human immunodeficiency virus type-1 (HIV-1) glycoprotein gp120. *Neuropharmacology* 2003;45:96-105.
91. Sanna MD, Peroni D, Quattrone A, Ghelardini C, Galeotti N. Spinal RyR2 pathway regulated by the RNA-binding protein HuD induces pain hypersensitivity in antiretroviral neuropathy. *Experimental neurology* 2015;267:53-63.
92. Kann O, Kovacs R. Mitochondria and neuronal activity. *Am J Physiol Cell Physiol* 2007;292:C641-57.
93. Douda DN, Khan MA, Grasemann H, Palaniyar N. SK3 channel and mitochondrial ROS mediate NADPH oxidase-independent NETosis induced by calcium influx. *Proceedings of the National Academy of Sciences of the United States of America* 2015;112:2817-22.

-
94. Korbecki J, Baranowska-Bosiacka I, Gutowska I, Chlubek D. The effect of reactive oxygen species on the synthesis of prostanoids from arachidonic acid. *J Physiol Pharmacol* 2013;64:409-21.
95. Gu X, Bo J, Zhang W, Sun X, Zhang J, Yang Y, Ma Z. Intrathecal administration of cyclic AMP response element-binding protein-antisense oligonucleotide attenuates neuropathic pain after peripheral nerve injury and decreases the expression of N-methyl-D-aspartic receptors in mice. *Oncol Rep* 2013;30:391-8.
96. Athos J, Impey S, Pineda VV, Chen X, Storm DR. Hippocampal CRE-mediated gene expression is required for contextual memory formation. *Nature neuroscience* 2002;5:1119-20.
97. Zhang JW, Klemm DJ, Vinson C, Lane MD. Role of CREB in transcriptional regulation of CCAAT/enhancer-binding protein beta gene during adipogenesis. *The Journal of biological chemistry* 2004;279:4471-8.