

Perioperative Opioid Usage Monitoring and Waste

Renyu Liu, MD, PhD* , John Grothusen, PhD and Scott Falk, MD

Department of Anesthesiology and Critical Care, Perelman School of Medicine at the University of Pennsylvania, USA

Abstract

This editorial discusses the status and issues related to perioperative opioid usage monitoring and waste. Opioid detection of wasted material is briefly discussed also. Flowlytics® from Invistics is a digital system to monitor opioid usage and waste in medical facilities. Opioid waste in medical facilities has a two-person witness procedure. Easy to use detection of wasted materials needs to be developed in the future. It is unclear whether the strategies used in medical facilities should be recommended for opioid disposal in the public to reduce opioid diversion. Relevant studies are needed.

Keywords

Opioid, Perioperative, Waste, Diversion, Monitoring, Detection

Introduction

The perioperative period is one of the places in a hospital where opioids are used on a daily basis. Areas where opioid diversion can occur in the hospital include the perioperative area where much of the diversion might go undetected. An early study published in 1987 in The Journal of the American Medical Association indicated that general practice, family medicine, and anesthesia are practice groups on the top of the list with high risk of substance dependence [1]. Anesthesia providers are more likely to have opioid use disorder (OUD) than other sub-specialties, with most of the abuse occurring through the intravenous route [1]. Anesthesiologists constitute 13-15% of people receiving substance dependence treatment in centers specialized in substance dependence management and monitoring for physicians [2]. One of the cited reasons why anesthesia providers have a higher incidence of OUD is their easy access to such medications [3]. Opioid usage for patients in hospital settings is generally very tightly controlled and monitored. Disposal of opioid waste in the perioperative area requires a two-person witness procedure, to ensure there are no discrepancies in the amount used in patients and the amount to be discarded. Disposal of opioids was less regulated in the past with unused opioids being dumped into regular trash or into a sink. Many practices now use a specific process bin or container to discard opioid medication.

Opioid usage for patients after surgery is generally managed by the surgical team. To improve the safety of opioid usage and deter diversion in the perioperative period, innovative initiatives and strategies need to be developed, studied and implemented.

Diversion Monitoring and Detection

Flowlytics® from Invistics was developed using artificial intelligence (AI) to monitor opioid usage and to detect potential diversion. A recent study indicated that use of machine learning and advanced analytics can detect known diversion cases much faster (ranges from 7-579 days faster) than traditional diversion detection methods that use periodic reporting [4]. This new methodology has demonstrated more than 95% accuracy, specificity, and sensitivity, and is designed to alert investigators for potential drug diversion by quickly detecting: 1) **Lack of or discrepancy in** drug reconciliation, 2) **Time-range matrix** to detect significant differences in drug waste behavior from a particular provider as compared to peers in similar cases and 3) A pattern of **full dose wasting** where a full vial of unused medication goes into waste instead of being returned [5]. The Flowlytics software has been part of the Senti 7 Suite from Wolters Kluwer since June 2023. It is a requirement now that opioid waste must be witnessed by a second person. In China, the opioid waste process is often video recorded, and all opioid ampoules must be returned to the pharmacy and documented. Figure 1 shows the artificial intelligence (AI) driven opioid waste and monitoring system developed by Rehn Meditech, China, for its patient controlled analgesia system, that ensures that opioid containing solutions are discarded properly and recorded. A recent study using hospital registries indicated that use of smaller syringes could reduce perioperative waste, cost and perioperative adverse effects [6]. Further research is needed to demonstrate whether such monitoring or data registry will reduce opioid diversion in the hospital settings.

Opioid Disposal in Perioperative Area

Opioids are the most commonly used pain medications in the perioperative period. Previously it was routine practice that unused opioids were dumped into trash bins or into a sink. Most opioids




Figure 1: The patient controlled analgesia medication unused medication returning system developed by Rehn Meditech (Courtesy of Rehn Meditech).



Figure 2: Opioid disposal container from Stericycle.

are, in fact, on the safe flush list of the USA Food and Drug Administration (FDA) [7]. In a study published in 2017, where 15 active ingredients were investigated for potential environmental impact, it was concluded that most of the investigated drugs flushed down the toilet do not pose significant environmental risk. However,

the authors did claim that additional data are needed [7]. With the increasing consumption of opioids, and the current opioid crisis, there is a growing concern of discarded opioids getting into the environment and generating negative impact on human and animal health. The guidelines on how to dispose opioids safely have

been updated in multiple safety agencies. The last update on the FDA website for “safe disposal of medicines” was November 2021 [8]. A recent study indicated that trace amounts of tramadol, a relatively weak opioid, in water could change fish behavior. Studies analyzing potent opioids, including morphine and heroin, in the wastewater of two USA cities with known drug overdose problems indicated that the number of overdose deaths correlated with the concentration of opioids detected in the wastewater [9]. To reduce the release of opioids into the environment, and deter potential diversion, new products for opioid disposal have been developed. Many institutions have adopted a policy that all unused opioids must be disposed of in a dedicated deactivation container. An example of a deactivation container from Stericycle is shown in Figure 2. The controlled substance is deactivated with activated charcoal with a bittering additive and solidifier added to further deter diversion.

Opioid Detection

Opioid detection is critical for not only patient and medical safety, but for environmental safety monitoring and national security. An ideal opioid detection system must have high sensitivity and specificity, and be easy to use with portability and a fast detection process that can be used in various field situations. Fentanyl strips are available now and are easy to use; however, they suffer sensitivity issues and false positive problems [10]. By using opioid receptor protein and graphene enabled technology, we and others have demonstrated high sensitivity in detecting opioids [11-13]. Such opioid protein based technology may ultimately prove to be the most sensitive and specific way to detect opioids and metabolites in the environment [12-15]. However, portability of such technology is limited at the present time.

In summary, this paper discusses the status of perioperative opioid monitoring and waste strategies to reduce potential diversion and OUD. It is unclear whether some of the strategies (for example, two-person witness procedure for opioid waste) will be suitable and adaptable for general public and household opioid management to reduce diversion and help to abate the OUD crisis. This topic is beyond the focus of this editorial. Further discussion and studies are needed before any new policy or guideline can be recommended.

Conflict Interest Statement

The authors has no relationship with any commercial product mentioned in this editorial. Dr. Renyu Liu is one of the inventors for the patents for water soluble mu opioid receptors (Patent number: 9670264), and the associated technologies to use water soluble opioid receptors to detect opioids (Patent number: US10809222B2).

Funding Support

Dr. Renyu Liu acknowledges the funding support from NIH for the water soluble mu opioid receptor related work (1R01GM111421, PI, RYL).

References

1. Talbott GD, Gallegos KV, Wilson PO, Porter TL. The Medical Association of Georgia's Impaired Physicians Program. Review of the first 1000 physicians: Analysis of specialty. *JAMA* 1987; 257: 2927-30.
2. Oreskovich MR, Caldeiro RM. Anesthesiologists recovering from chemical dependency: Can they safely return to the operating room? *Mayo Clin Proc* 2009; 84: 576-80.
3. Misra U, Gilvarry E, Marshall J, Hall R, McLure H, Mayall R, et al. Substance use disorder in the anaesthetist: Guidelines from the Association of Anaesthetists. *Anaesthesia* 2022; 77: 691-699.
4. Knight T, May B, Tyson D, McAuley S, Letzkus P, et al. Detecting drug diversion in health-system data using machine learning and advanced analytics. *Am J Health Syst Pharm* 2022; 79: 1345-1354.
5. Solutions R. RXpert Solutions - An inside look Flowlytics with Xing Guan, 2022. <https://youtu.be/QuEGeg61onk?si=dLTWljqmKgDxjHQz>
6. Redaelli S, Suleiman A, von Wedel D, Ashrafian S, Munoz-Acuna R, et al. Intraoperative opioid waste and association of intraoperative opioid dose with postoperative adverse outcomes: A hospital registry study. *Pain Ther* 2024; 13: 211-225.
7. Khan U, Bloom RA, Nicell JA, Laurenson JP. Risks associated with the environmental release of pharmaceuticals on the U.S. Food and Drug Administration "flush list". *Sci Total Environ* 2017; 609: 1023-1040.
8. FDA: Safe Disposal of Medicines, 2021. <https://www.fda.gov/drugs/buying-using-medicine-safely/safe-disposal-medicines>
9. Gushgari AJ, Venkatesan AK, Chen J, Steele JC, Halden RU. Long-term tracking of opioid consumption in two United States cities using wastewater-based epidemiology approach. *Water Res* 2019; 161: 171-180.
10. Halifax JC, Lim L, Ciccarone D, Lynch KL. Testing the test strips: laboratory performance of fentanyl test strips. *Harm Reduct J* 2024; 21: 14.
11. Kumar N, Rana M, Geiwitz M, Khan NI, Catalano M, et al. Rapid, Multianalyte Detection of Opioid Metabolites in Wastewater. *ACS Nano* 2022; 16: 3704-3714.
12. Lerner MB, Matsunaga F, Han GH, Hong SJ, Xi J, et al. Scalable production of highly sensitive nanosensors based on graphene functionalized with a designed G protein-coupled receptor. *Nano Lett* 2014; 14: 2709-14.
13. Liu R, Johnson ATC, Saven JG. Water soluble G-protein coupled receptor enabled biosensors. *Transl Perioper Pain Med* 2019; 6: 98-103.
14. De-Eknamkul C, Zhang X, Zhao MQ, Huang W, Liu R, et al. MoS(2)-enabled dual-mode optoelectronic biosensor using a water soluble variant of mu-opioid receptor for opioid peptide detection. *2d Mater* 2020; 7(1): 014004. doi: 10.1088/2053-1583/ab5ae2
15. Wen C, Selling B, Yeliseev A, Xi J, Perez-Aguilar JM, et al. The C-terminus of the mu opioid receptor is critical in G-protein interaction as demonstrated by a novel graphene biosensor. *IEEE Sens J* 2021; 21: 5758-5762.

Citation: Liu R, Grothusen J, Falk S. Perioperative Opioid Usage Monitoring and Waste. *Transl Perioper Pain Med* 2024; 11(1):597-600

Copyright: © 2024 Liu R, et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

***Corresponding Author:** *Renyu Liu, MD, PhD, Professor, Departments of Anesthesiology and Critical Care, and Neurology, Perelman School of Medicine at the University of Pennsylvania. 336 John Morgan building, 3620 Hamilton Walk, Philadelphia, PA 19104, USA, E-mail: RenYu.Liu@pennmedicine.upenn.edu*

Editor: *Li-Ming Zhang, MD, Associate Professor of Anesthesiology, Attending Anesthesiologist (Recertified), UPP Dept of Anesthesiology and Perioperative Medicine, 200 Lothrop Street, UPMC MUH Suite N-467, Pittsburgh, PA 15213, USA; Tel: 412-648-6077; Fax: 412-648-6014; Lab: 412-624-8386; E-mail: zhangl1@anes.upmc.edu*

Additional publication details

Journal short name: *Transl Perioper Pain Med*

Received Date: April 23, 2024

Accepted Date: May 02, 2024

Published Date: May 07, 2024